



Letters to the Editor

COVID-19 and the worsening of health inequities in Santiago, Chile

Usama Bilal ,^{1,2,*} Tania Alfaro³ and Alejandra Vives^{4,5}

¹Urban Health Collaborative, Drexel Dornsife School of Public Health, Philadelphia, PA, USA, ²Department of Epidemiology and Biostatistics, Drexel Dornsife School of Public Health, Philadelphia, PA, USA, ³Escuela de Salud Pública, Universidad de Chile, Santiago, Chile, ⁴Departamento de Salud Pública, Pontificia Universidad Católica de Chile, Santiago de Chile, Chile and ⁵Centro de Desarrollo Urbano Sustentable (CEDEUS), Santiago de Chile, Chile

*Corresponding author. 3600 Market St, Suite 730, Philadelphia, PA, 19104, USA. E-mail: ubilal@drexel.edu

Editorial decision 3 January 2021; Accepted 8 January 2021

Key words: Mortality, health equity, inequalities, urban health, Latin America, Chile

The COVID-19 pandemic is already responsible for >1 million deaths worldwide.¹ Latin America is one of the most affected regions worldwide, with >300 000 deaths confirmed by September 2020.¹ Latin America is also one of the most urbanized and unequal regions in the world, with wide inequities in longevity in its cities.² Wide inequities in COVID-19 outcomes have been reported in other settings.³ However, policymakers in some Latin American countries have expressed scepticism about the existence of health inequities in COVID-19 mortality.⁴ We used mortality, population and census data to show a worsening of pre-existing inequities in mortality in the municipalities that make up the metropolitan region of Santiago (Chile) during the COVID-19 pandemic.

We obtained data for the 36 *comunas* (municipalities) that make up Greater Santiago, the metropolitan area of the capital of Chile, on: (i) mortality by age for the 2016–2020 period from the Department of Statistics and Health Information (DEIS); (ii) population projections by age for the 2016–2020 period from the National Institute of Statistics (INE); and (iii) average years of schooling among adults aged ≥25 years, and the proportion of households living in overcrowding (>2.5 people/bedroom) from the 2017 Chilean census. We selected these two indicators since they represent either good measures of area-level socio-economic status (SES) or are directly linked to COVID-19 transmission. We computed age-adjusted mortality rates, using the 2000 World Health Organization reference population, from January to August for the pre-pandemic (years 2016–2019) and pandemic (2020) periods. We estimated the association between log(mortality) and average years of schooling or proportion overcrowded households using a linear model for each period separately. Data and code for replication are available here: https://github.com/usamabilal/COVID_Chile_Inequities.

Figure 1 shows the main results. We found a strong association between SES and mortality in both periods, although this association was stronger in 2020. Specifically, we found a 9.0% lower mortality per 1-year increase in the average

schooling years in the pre-pandemic period [relative risk (RR) = 0.91, 95% confidence interval (CI) 0.87 to 0.93] compared with a 13.8% lower mortality in the 2020 period (RR = 0.86, 95% CI 0.83 to 0.89). We also found that a 5% increase in the proportion of overcrowded households was associated with a 22% and 32% higher mortality in the pre-pandemic and pandemic periods, respectively (RR = 1.22, 95% CI 1.16 to 1.28; RR = 1.32, 95% CI 1.23 to 1.42).

In summary, we found a worsening of pre-existing inequities in mortality in the metropolitan area of Santiago de Chile during the COVID-19 pandemic. The association of years of schooling and overcrowding with mortality in the pre-pandemic period (2016–2019) is consistent with previous research in Santiago, showing wide gaps in life expectancy² and infant mortality.⁵ Our finding that mortality inequities in 2020 were greater than in previous periods contradicts statements made in September 2020 by the Minister of Health of Chile, which indicated that there is no relationship between mortality due to COVID-19 and poverty, as this would imply discriminatory healthcare due to the place of origin of the patients.⁴ This assertion ignores existing structural inequalities in Chile, including inequities in healthcare access and utilization by type of insurance both before⁶ and during the pandemic,⁷ along with social determinants of health beyond the health system. These include factors driving not only incidence and increased exposure to the virus,³ such as more precarious employment conditions, loss of income and the need to work outside the home, but also an increased prevalence of conditions that aggravate the consequences of the disease, which show a strong social patterning in Chile.⁸ Mitigating health inequities was an explicit objective of the Chilean National Health Strategy⁹ and the aggravation of these inequities amidst the COVID-19 pandemic represents a step in the wrong direction. Our analysis is limited by the lack of individual-level mortality data and a validated SES index. However, we are using whole-




Figure 1 Area-level socio-economic status and age-adjusted mortality in 2016–2019 and 2020 in the municipalities of Santiago, Chile (A) Average years of schooling for adults >25 years of age; (B) percentage of households living in overcrowded situations. The solid line represents a linear fit.

population vital-registration data in a country with a good vital-registration system,¹⁰ which lends strength to our findings.

Interventions to reduce these inequities are greatly needed to avoid the continued widening of these gaps, including but not exclusively focused on healthcare, in addition to addressing the other health and economic consequences of the pandemic, which will likely also affect these vulnerable groups more frequently, intensely and for more prolonged periods of time.

Funding

U.B. was supported by the Office of the Director of the National Institutes of Health under award number DP5OD26429. U.B., T.A. and A.V. were also supported by the Salud Urbana en América Latina (SALURBAL)/Urban Health in Latin America project, funded by the Wellcome Trust (205177/Z/16/Z). The funding sources had no role in the analysis, writing or decision to submit the manuscript.

Acknowledgements

The data underlying this article are available at https://github.com/usamabil/COVID_Chile_Inequities/. All analyses were conducted with publicly available de-identified surveillance data.

Conflict of interest

None declared.

References

1. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis* 2020;20:533–34.
2. Bilal U, Alazraqui M, Caiaffa WT *et al*. Inequalities in life expectancy in six large Latin American cities from the SALURBAL study: an ecological analysis. *Lancet Planet Health* 2019;3:e503–10.
3. Berkowitz RL, Gao X, Michaels EK, Mujahid MS. Structurally vulnerable neighbourhood environments and racial/ethnic COVID-19 inequities. *Cities & Health* 2020;1–4.
4. Basoalto H, Silva D. Ministro Paris por fallecidos por Covid: ‘No hay una relación entre letalidad y pobreza, eso significaría que los pacientes son atendidos en forma discriminada’. *La Tercera* 2020. <https://www.latercera.com/nacional/noticia/ministro-paris-por-fallecidos-por-covid-no-hay-una-relacion-entre-letalidad-y-pobreza-eso-significaria-que-los-pacientes-son-atendidos-en-forma-discriminada/S5LPFNI4VRDZ7EP7FHJJEVJNJ2A/>

5. Gonzalez F, Alfaro T, Horna-Campos O. Mortalidad Infantil en las comunas de la Región Metropolitana, periodo 2005-2014. *Rev Chil Pediatr* 2020;**91**:34–45.
6. Castillo-Laborde C, Aguilera-Sanhueza X, Hirmas-Adaury M *et al.* Health insurance scheme performance and effects on health and health inequalities in Chile. *MEDICC Review* 2017;**19**:57–64.
7. Escuela de Salud Pública de la Universidad de Chile, Colegio Médico de Chile. *Monitoreo Nacional de Síntomas y Prácticas COVID-19 en Chile (MOVID-19): ¿Cuál ha sido el impacto de la pandemia en el acceso a atenciones de salud? Un análisis para la adaptación de nuestro sistema de salud.* 2020. <https://www.movid19.cl/publicaciones/decimo-informe/>
8. Ortiz MS, Cabieses B, Oyarte M, Repetto P. Disentangling socioeconomic inequalities of type 2 diabetes mellitus in Chile: a population-based analysis. *PLoS One* 2020;**15**:e0238534.
9. Ministerio de Salud (Chile). *Estrategia Nacional de Salud.* Gobierno de Chile Town: Santiago de Chile. 2011.
10. Bilal U, Hessel P, Perez-Ferrer C *et al.* Life expectancy and mortality in 363 cities of Latin America. *Nature Medicine* 2021. <https://doi.org/10.1038/s41591-020-01214-4>

BMI is unlikely to be a plausible intervention target for reducing the incidence of dementia

International Journal of Epidemiology, 2021, 1040–1041

doi: 10.1093/ije/dyab062

Advance Access Publication Date: 13 April 2021



Neil M Davies ^{1,2,3,*} Roxanna Korologou-Linden^{1,2} and Emma L Anderson^{1,2}

¹Medical Research Council Integrative Epidemiology Unit, University of Bristol, Bristol, UK, ²Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK and ³K.G. Jebsen Center for Genetic Epidemiology, Department of Public Health and Nursing, NTNU, Norwegian University of Science and Technology, Norway

*Corresponding author. Population Health Sciences, Bristol Medical School, University of Bristol, Barley House, Oakfield Grove, Bristol, BS8 2BN, UK. E-mail: Neil.Davies@bristol.ac.uk

Editorial decision 3 January 2021; Accepted 8 January 2021

Ma and colleagues investigated the relationship between body mass index (BMI) in midlife ($n = 6582$; average age, 62.5 years) and risk of dementia using data from the English Longitudinal Study of Ageing. They report that those who were obese had a 34% [95% confidence interval (CI), 7–61%] increased risk of being diagnosed with dementia over a 15-year follow-up of the study compared with those who were non-obese at baseline.¹ Associations were adjusted for a range of baseline covariates including age, sex, apolipoprotein E- $\epsilon 4$ allele status, education, marital status, current smoking, physical activities, hypertension and diabetes. The positive relationship reported by Ma and colleagues is surprising in light of previous observational studies, including one of 2 million participants suggesting the opposite direction of association.²

Given these results, is it likely that obesity is a plausible modifiable risk factor for dementia? There have been many studies showing that obesity in midlife is associated with risk of dementia. However, as we know, correlation does not equal causation. The authors correctly note that the incidence of obesity has increased in recent years and, if their estimates did indeed reflect the true causal effect of obesity on risk of dementia, then rates of dementia could be expected to increase in the future. However, some of the best available evidence to date suggests that the age-specific incidence of Alzheimer's disease has declined over time in the UK.³

To definitively prove that obesity increases the risk of dementia, we would ideally need to conduct a large randomized-controlled trial of an intervention that modifies BMI (e.g. bariatric surgery). The intervention would have to occur long before the expected onset of dementia,⁴ as most types of dementia have a long prodromal

period. We would then need to follow up those participants over time to see whether those who received the surgery did indeed have a lower risk of dementia. However, this hypothetical trial would be expensive, time-consuming and potentially unethical to conduct.

An analogous approach is to exploit naturally occurring random variation in BMI as a proxy. For example, genetic variation is known to be robustly associated with, and explains ~6% of the variation in BMI including variants near the FTO gene on chromosome 16.⁵ If BMI affects risks of dementia, then we would expect individuals with genetic variants that predispose them to higher levels of BMI across the life course to also have a higher risk of dementia and the most common type of dementia, Alzheimer's disease. This approach, known as Mendelian randomization (MR), is less likely to suffer from bias due to confounding and cannot be affected by reverse causation, because genetic variants are randomly allocated at conception and cannot be modified by subsequent disease. MR estimates suggest that increases in BMI are unlikely to have any clinically meaningful effect on Alzheimer's disease (odds ratio per unit increase in BMI: 0.98; 95% CI, 0.89–1.08 in Korologou-Linden and colleagues; and odds ratio: 1.05; 95% CI, 0.91–1.21 in Larsson and colleagues).^{6,7} However, there is evidence that genetic liability to Alzheimer's disease has a causal effect on BMI and the magnitude of this effect increases with age, suggesting that the associations of BMI and Alzheimer's disease reported by observational studies are likely a result of reverse causation or survival bias.

As with all research methods, MR has limitations and can suffer from selection or collider bias, or survival effects. MR assumes a linear effect of the risk factor and may not be able to detect U-shaped